

界面活性劑의 合成技術


공학박사 남기대 교수
충북대학교 공업화학과

3 Synthesis of Surfactants

Dr. M. Biermann, Henkel KGaA, D-4000 Düsseldorf
Dr. F. Lange, Henkel KGaA, D-4000 Düsseldorf
Dr. R. Piott, Henkel KGaA, D-4000 Düsseldorf
Dr. U. Ploog, Henkel KGaA, D-4000 Düsseldorf
Dr. H. Rutzen, Henkel KGaA, D-4000 Düsseldorf
Dr. J. Schindler, Henkel KGaA, D-4000 Düsseldorf
Dr. R. Schmid, Henkel KGaA, D-4000 Düsseldorf

3.1	Raw Materials
3.1.1	Fat Derived Chemical Raw Materials
3.1.1.1	Fatty Acids
3.1.1.2	Fatty Acid Methyl Esters
3.1.1.3	Fatty Alcohols
3.1.1.4	Fatty Amines
3.1.2	Petrochemical Raw Materials
3.1.2.1	Long-Chain Hydrocarbons
3.1.2.2	Short-Chain Hydrocarbons
3.1.2.3	Alkylene Oxides
3.1.2.4	Olefins, Fatty- and Oxo-Alcohols – by Chain-Building Reactions
3.1.2.5	Alkyl Aromatics
3.1.2.6	Amines
3.1.3	Fatty Chemical and Petrochemical Raw Materials – Contrast or Supplement?
3.2	Surfactants
3.2.1	Anionic Surfactants
3.2.1.1	General Production Technologies
3.2.1.2	Soaps
3.2.1.3	Alkylbenzene Sulfonates
3.2.1.4	Fatty Alcohol Sulfates
3.2.1.5	Alkyl Ether Sulfates
3.2.1.6	Alkane Sulfonates
3.2.1.7	Olefin Sulfonates
3.2.1.8	α -Sulfo Fatty Acid Esters
3.2.1.9	Internal Sulfo Fatty Acid Esters
3.2.1.10	Fatty Acid Cyanamides (Amide Soaps)
3.2.1.11	Sulfo Succinic Acid Alkyl Esters
3.2.1.12	Acyl Oxyalkane Sulfonates (Fatty Acid Isethionates)
3.2.1.13	Acylamino Alkane Sulfonates (Taurides)
3.2.1.14	Ether Sulfonates

3.2.1.15	Ether Carboxylic Acids
3.2.1.16	Sarcosinates
3.2.1.17	Alkyl Phosphates and Alkyl Ether Phosphates
3.2.2	Nonionic Surfactants
3.2.2.1	Basic Properties
3.2.2.2	Polyglycol Ethers
	Fatty alcohol polyglycol ethers
	Alkylphenol polyglycol ethers
	End group-capped fatty alcohol polyglycol ethers
	Fatty acid polyglycol esters
	Fatty acid alkanol amides, fatty acid alka.olamide polyglycol ethers
	Ethylene oxide/propylene oxide block polymers
3.2.2.3	Polyol Surfactants
	Glycerine fatty acid esters
	Sorbitan esters
	Sugar derived surfactants (sugar esters, alkyl glycosides).
3.2.2.4	Amine Oxides
3.2.3	Cationic Surfactants
3.2.3.1	Application Areas
3.2.3.2	Reaction Conditions
	Amine salts
	Quaternary ammonium compounds (quats)
3.2.3.3	Amines with one long Alkyl Group
3.2.3.4	Amines with a long β -Hydroxyalkyl Group
3.2.3.5	Amines with Several long Alkyl Groups
3.2.3.6	Amines with Several long β -Hydroxyalkyl Groups
3.2.4	Amphoeritic Surfactants
3.2.4.1	Carboxy Amphoeritic Surfactants
	Alkyl betaines
	Other amphoteric
3.2.4.2	Sulfo Betaines
3.2.4.3	Basic Properties
3.2.5	Biosurfactants
3.2.5.1	Structure and Synthesis
3.2.5.2	Applications
3.3	Literature



※ 계면활성제의 새로운 문제점

- 1) 정교한 기능을 갖는 고부가가치성의 요구
- 2) 환경오염 방지책 - 천연에서 유도된 새로운 조성을 갖어 생분해가 좋은 것
- 3) 새로운 조성과 기능을 갖는 화합물 개발

※ 새로운 조성을 갖는 계면활성제 창안

당류를 친수성기로한 비이온성계면활성제

알파솔폰고급지방산염인 음이온성계면활성제

에스테르화

당류 및 솔픈산기를 갖는 고급지방산 에스테르형

※ 새로이 합성된 화합물 기대성

- 1) 생물학적 특이성 갖는 세정성
- 2) 도료, 인쇄, 농약의 특이한 분산성
- 3) 섬유의 대전방지성
- 4) 고분자 합성에 있어 새로운 유화제 개발


Synthesis of Polyolic α -Sulfo Long Chain Alcanoates.

Materials ; 1) α -sulfo fatty acid - Hydrocarbon No. 12~18

2) Sugar ; glycerol, *myo*-inositol


D-mannitol, sorbitol

1) 1-mono glycerol mono α -sulfo alkanate




glycerol


1,2-O-isopropylidene glycerine
 ● remove H₂O of azotropic distillation with benzene



2) diglycerol mono α -sulfo alkanoates





dichloro methane
boron trifluoride-ether complex




diglycerol mono α -sulfo alkanoic acid

3) Symmetrical triglycerol mono α -sulfo alkanoate





2,2-dimethyl-1,3-dioxolane
→ methanol

1,3-dichloro-2-propanol




diisopropyliden triglycerol




Symmetrical diisopropyliden triglycerol mono α -sulfo alkanoates

borotrifluoride-ether complex
dichloromethane



Symmetrical triglycerol mono α -sulfo alkanoate

메틸 α -D-글루코파라노사이드 유도체




Scheme The starting materials, intermediates and reacted products.


R : C₉H₁₉-, C₁₁H₂₃-, C₁₃H₂₇-, C₁₅H₃₁-, C₁₇H₃₅-

$$X + Y + Z = 20$$


계면활성 아미노 당 유도체




- [I] R = decyl
- [II] R = dodecyl
- [III] R = tetradecyl
- [IV] R = hexadecyl




- [V] R = octyl
- [VI] R = decyl
- [VII] R = dodecyl
- [VIII] R = tetradecyl



- [VI] R = octyl
- [VII] R = decyl
- [VIII] R = dodecyl
- [VIII] R = tetradecyl




- [XIII] R = octyl
- [XIV] R = decyl
- [XV] R = dodecyl
- [XVI] R = tetradecyl




- [XVII] R = octyl
- [XVIII] R = decyl
- [XIX] R = dodecyl
- [XX] R = tetradecyl



I. Polyolic Nonionic Surfactant

1) Myo-inositol Monoester





4) myo-inositol mono α -sulfo alkanoate






hydrolysis of saturated methanolic ammon



myo-inositol-2-mono α -sulfo alkanoate

5) Esterification of α -sulfo alcanoate of D-mannitol or Sorbitol



Synthesis method

① Direct esterification

solvent ; chloroform, benzene, toluene


azotropic removal of water

② Acid chloride method

solvent ; dimethylformamide(DMF) + pyridine

α -sulfo alcanoic acid dissolved with CCl_4

③ Alcoholysis
















solvent ; DMF, benzene









catalyst ; MeONa

azotropic removal of methanol

reaction apparatus - Fig. 10


베타-인 헝 계면활성제






N-일킬 카르복시 베타인 유도체


Scheme The process for the synthesis of N-alkylcarboxybetaines.



- (1a) : N-(2-decanoxyethyl)-N,N-dimethylammonio ethanoate
- (1b) : N-(2-dodecanoxyethyl)-N,N-dimethylammonio ethanoate
- (1c) : N-(2-tetradecanoxyethyl)-N,N-dimethylammonio ethanoate
- (1d) : N-(2-hexadecanoxyethyl)-N,N-dimethylammonio ethanoate



- (2a) : N-(2-decanoxyethyl)-N,N-dimethylammonio ethanoate
- (2b) : N-(2-dodecanoxyethyl)-N,N-dimethylammonio ethanoate
- (2c) : N-(2-tetradecanoxyethyl)-N,N-dimethylammonio ethanoate
- (2d) : N-(2-hexadecanoxyethyl)-N,N-dimethylammonio ethanoate




(3a) : N-(2-decanoylaminoethyl)-N,N-dimethylammonio ethanoate

(3b) : N-(2-dodecanoylaminoethyl)-N,N-dimethylammonio ethanoate







(3c) : N-(2-tetradecanoylaminoethyl)-N,N-dimethylammonio ethanoate

(3d) : N-(2-hexadecanoylaminoethyl)-N,N-dimethylammonio ethanoate


이미다졸린 형 계면활성제






1-(2-hydroxyethyl)- 2-alkyl-2-imidazoline HEAI




VIII.



IX.



X.





XI.




II. Maleic Anhydride - Polyethylene Glycol-Phthalic Anhydride
Polymeric Surfactant


Step 1.



Step 2.






Step 3.






1. Anionic Polymeric Surfactant

1. Vinyl α -Sulfo Fatty Acid Oligomer




R : C₁₀ ~ C₁₆


2. Allyl α -Sulfo Long Chain Aliphatic Carboxylate Polymers




3. 알파술운지 방산 음이온성 울리고며 계면활성제



(R = C₈, C₁₀, C₁₂, C₁₄, C₁₆)




(n = 5, 10, 20 , R' = C₁₂H₂₅)





alkyl polyoxyethylene α -sulfo alkanoate








모노 글리세레이트류 유도체






모노 알킬 인산염(MAP)류 유도체





슬포 속신산 염류 유도체




메틸 타우린산 염류 유도체





N-아실 이세티온산 염류 유도체



아실화 콜라겐 폴리펩티드류 유도체










아미노산 모노 글리세리드 에스테르류
















N-알킬 글루타민산 염류 유도체

I. Schotten - Baumann 법













■. Silicon Surfactant


1. Poly Siloxane Nonionic Surfactant

a) ω -Hydroxylated polysiloxane




n : 2, 3, 4


b) Acetoxy pentamethyl disiloxane




c) ω -oxy pentamethyl disiloxane









d) Bis (pentamethyl disiloxyl) methyl silane



e) Tris (trimethyl siloxy) silane





f) Silicon Nonionic Surfactant




N. Polyolic Silicon Surfactant

1. Polyoxy Siloxane





2. Allyl Glycidil Alcohol




b) Sucrose, c) Glucose, d) Glycerine

3. Polyolic Silicon Surfactant



1) Polyolic primary polymethyl siloxane(A)



[1a] ; m=2, n=6.5 P-2-MEO-6.5

[1b] ; m=3, n=6.5 P-3-MEO-6.5


[1c] ; m=4, n=6.5 P-4-MEO-6.5

[1d] ; m=2, n=12 P-2-MEO-12

[1e] ; m=3, n=12 P-3-MEO-12

[1f] ; m=4, n=12 P-4-MEO-12

2) Polyolic secondary polymethyl siloxane(B)



[2a] ; m=2, n=6.5 S-2-MEO-6.5

[2b] ; m=3, n=6.5 S-3-MEO-6.5


[2c] ; m=4, n=6.5 S-4-MEO-6.5

[2d] ; m=2, n=12 S-2-MEO-12

[2e] ; m=3, n=12 S-3-MEO-12

[2f] ; m=4, n=12 S-4-MEO-12

3) Polyolic tertiary polymethyl siloxane(C)



[3a] ; m=2, n=6.5 T-2-MEO-6.5

[3b] ; m=3, n=6.5 T-3-MEO-6.5


[3c] ; m=4, n=6.5 T-4-MEO-6.5

[3d] ; m=2, n=12 T-2-MEO-12

[3e] ; m=3, n=12 T-3-MEO-12


[3f] ; m=4, n=12 T-4-MEO-12

4) Polyolic side-chain modified polymethyl siloxane(D)



[4a] ; m=2, n=6.5 PH-2-MEO-6.5

[4b] ; m=6, n=6.5 PH-6-MEO-6.5



[5a] ; m=2, n=6.5 D-2-PH-2-MEO-6.5

[5b] ; m=4, n=6.5 D-2-PH-4-MEO-6.5

